

An almost isotropic cosmic microwave temperature does not imply an almost isotropic universe

U. S. Nilsson

*Department of Applied Mathematics
University of Waterloo
Waterloo, Ontario N2L 3G1, Canada*

C. Ugglä

*Department of Engineering Sciences, Physics and Mathematics
University of Karlstad
S-651 88 Karlstad, Sweden*

J. Wainwright

*Department of Applied Mathematics
University of Waterloo
Waterloo, Ontario N2L 3G1, Canada*

and

W. C. Lim

*Department of Applied Mathematics
University of Waterloo
Waterloo, Ontario N2L 3G1, Canada*

ABSTRACT

In this letter we will show that, contrary to what is widely believed, an almost isotropic cosmic microwave background (CMB) temperature does not imply that the universe is “close to a Friedmann-Lemaître universe”. There are two important manifestations of anisotropy in the geometry of the universe, (i) the anisotropy in the overall expansion, and (ii) the intrinsic anisotropy of the gravitational field, described by the Weyl curvature tensor, although the former usually receives more attention than the latter in the astrophysical literature. Here we consider a class of spatially homogeneous models for which the anisotropy of the CMB temperature is within the current observational limits but whose Weyl curvature is not negligible, i.e. these models are not close to isotropy even though the CMB temperature is almost isotropic.

Subject headings: Cosmology: theory – cosmic microwave background

Observations indicate that the temperature of the CMB is isotropic to a remarkable degree. This observation is widely believed to indicate that the universe is almost isotropic in the sense that it is “close to a Friedmann-Lemaître (FL) model”, at least since the time of last scattering. Support for this belief is provided by a fundamental result of Ehlers, Geren & Sachs (1968), known as the EGS theorem, which states that if the CMB temperature were exactly isotropic about every point in spacetime, the universe would have to be exactly an FL model. This result is of course not directly applicable to the real universe since the CMB temperature is not exactly isotropic. It is thus of considerable interest to note that Stoeger *et al.* (1995) were able, by making what might seem to be reasonable assumptions, to extend the results of Ehlers *et al.* by replacing the word “exactly” by “almost”, thus obtaining an “almost” EGS theorem. More precisely, they showed, given certain hypotheses, that if the CMB temperature is measured to be almost isotropic in a spacetime region of an expanding universe, then the universe is close to an FL model *in that region*.

Before proceeding further, it is necessary to give a precise meaning to the phrase “close to an FL model”. We follow the approach given in Wainwright & Ellis (1997, section 2.4), but focus primarily on the question of anisotropy rather than inhomogeneity. To quantify the deviation of the universe from an FL model as regards to anisotropy, the shear tensor σ_{ab} (see Wainwright & Ellis 1997, page 18 for the definition) of the cosmological fluid and the Weyl curvature tensor C_{abcd} are of primary importance since both are zero in an FL model. The shear tensor describes anisotropy in the overall expansion of the universe, while the Weyl curvature tensor represents the “free gravitational field”, describing for example, tidal forces. It is related to time derivatives of the shear tensor (see Wainwright & Ellis 1997, Eqs. (1.43) and (1.47)). Requiring the above quantities to be small in an anisotropic ever-expanding cosmology is not sufficient since these quantities are not dimensionless. Moreover, the shear, for example, tends to zero at late times irrespective of whether the model isotropizes or not. The appropriate quantities are the dimensionless ratios formed by normalizing with the Hubble scalar H (see Wainwright & Ellis 1997, page 18 for the definition), thus measuring the dynamical importance of the different variables with respect to the overall expansion of the universe. This insight dates back

to the important paper on observations in cosmology by Kristian & Sachs (1966, see page 398), in which they showed that various geometrical quantities, including the Weyl curvature tensor, can in principle be restricted by observations of distant galaxies (see their Table 1 for a list of their so-called anisotropy and inhomogeneity parameters). We therefore introduce the parameters Σ and $\mathcal{W}$ according to,

$$\Sigma^2 = \frac{\sigma_{ab}\sigma^{ab}}{6H^2}, \quad \mathcal{W}^2 = \frac{E_{ab}E^{ab} + H_{ab}H^{ab}}{6H^4}, \quad (1)$$

where σ_{ab} is the shear tensor and E_{ab} and H_{ab} are the electric and magnetic parts of the Weyl curvature tensor respectively. They are defined by

$$E_{ac} = C_{abcd}u^b u^d, \quad H_{ac} = \frac{1}{2}\epsilon_{ab}{}^{ef}C_{efcd}u^b u^d. \quad (2)$$

where u^a is the 4-velocity of the cosmological fluid. We refer to Σ as the *shear parameter* and to $\mathcal{W}$ as the *Weyl parameter*. For non-tilted spatially homogeneous perfect fluid models, a zero shear tensor implies that the Weyl curvature tensor is zero, and thus characterizes the FL models. However, restricting the Hubble-normalized shear to be small does *not* guarantee that the Hubble-normalized Weyl curvature is small since derivatives of σ_{ab} need not be small compared to H^2 . Therefore a necessary condition for the universe to be close to an FL model is that the Hubble-normalized shear *and* the Hubble-normalized Weyl curvature are small. In terms of the previously introduced anisotropy parameters, this necessary condition becomes

$$\Sigma \ll 1, \quad \mathcal{W} \ll 1. \quad (3)$$

To characterize the anisotropies of the CMB temperature, a spherical harmonic expansion of the temperature of the CMB sky,

$$\frac{\Delta T}{T} = \frac{T(\theta, \varphi) - \bar{T}}{\bar{T}} = \sum a_{lm} Y_{lm}(\theta, \varphi), \quad (4)$$

is often used. In the above expression, $\bar{T}$ is the mean temperature of the CMB sky and (θ, φ) are the usual spherical angles. The multipoles,

$$a_l = \left(\sum_{m=-l}^l a_{lm}^2 \right)^{1/2}, \quad (5)$$

describe, coordinate independently, the deviation of the CMB temperature from perfect isotropy. The

dipole anisotropy, a_1 , is commonly interpreted as describing the motion of the solar system with respect to the rest frame of the CMB, see e.g. Lineweaver *et al.* (1996), making the quadrupole, a_2 , the lowest multipole to describe “true” anisotropies. The observational bounds on the quadrupole and octupole are of the order (Stoeger *et al.* 1997)

$$a_2 \approx 10^{-5}, \quad a_3 \approx 10^{-5}. \quad (6)$$

The line of research initiated by Stoeger *et al.* (1995) into the implications of CMB temperature observations has been extended by Maartens *et al.* in a series of papers (Maartens *et al.* 1995a, 1995b, 1996) to give explicit bounds on various anisotropy and inhomogeneity parameters. It turns out that to first order, only the first three multipoles: the dipole, quadrupole and octupole, play a direct role, via the Einstein-Boltzmann equations, in limiting the anisotropy and inhomogeneity parameters. Specifically, Maartens *et al.* (1995b) deduced general inequalities from which we infer the following upper bounds on the present values of the shear and Weyl parameters, using the observational bounds on a_2 and a_3 in Eq. (6):

$$\Sigma_o < 10^{-4}, \quad \mathcal{W}_o < 10^{-3}. \quad (7)$$

These follow from Eqs. (31), (35), and (36) in Maartens *et al.* (1995b) using our Eq. (1). It is important to note, however, that these upper bounds are valid only if a number of hypotheses about the behavior of the multipole moments in the universe since last scattering are true. The essential hypotheses can be stated as follows:

- H_1 : the Copernican principle holds approximately,
- H_2 : the multipole moments have never been larger than they are today,
- H_3 : the dimensionless time and spatial derivatives of the multipoles are bounded by the multipoles themselves (Maartens *et al.* 1995b, assumptions C1' and C2').

The observation of the CMB temperature multipoles at our present position and at the present time, in conjunction with hypotheses H_1 and H_2 , gives bounds on the multipoles in a spacetime region between last scattering and the present, encompassing all fundamental observers with whom we have been in causal

contact with since last scattering (Maartens *et al.* 1995a, Eqs. (42) and (44)). The derivatives of the multipoles are not directly accessible to us (Maartens *et al.* 1996, page L9) since to obtain this information would require observations over time and space intervals on cosmological scales. A close examination of the paper by Maartens *et al.* (1995b, Eqs. (24) - (29)) shows that all of the upper bounds on the various anisotropy and inhomogeneity parameters depend on hypothesis H_3 and are thus not, in practice, observationally testable. Thus, *if hypothesis H_3 is not satisfied then the CMB temperature observations*, in conjunction with the analysis of Maartens *et al.* (1995b), *do not impose upper bounds on Σ and $\mathcal{W}$* , and hence do not establish that the universe is close to FL. It is clearly of interest to ask whether one can construct model universes which illustrate this possible limitation of the CMB temperature observations.

In this letter we exhibit such a class of cosmological models, namely the spatially homogeneous non-tilted Bianchi type VII₀ dust models in which the CMB is treated as a test field.¹ To obtain the present CMB temperature pattern, the photon energies are integrated along the null geodesics that connect points of emission on the surface of last scattering with the event of observation at the present time. It will be assumed that the process of matter-radiation decoupling in the early universe took place instantaneously and that the CMB photons travel to us along null geodesics from a redshift of $z \approx 1100$. If T_e is the temperature of the CMB at the surface of last scattering, the observed temperature T_o is given by (Nilsson *et al.* 1998)

$$T_o = T_e \exp \left\{ - \int_{\tau_e}^{\tau_o} [1 + \Sigma_{ab}(\tau) K^a(\tau) K^b(\tau)] d\tau \right\}, \quad (8)$$

where the subscripts o and e refer to observation and emission respectively. Here K^a are the directional cosines of a particular photon path and $\Sigma_{ab} = \sigma_{ab}/H$ is the Hubble-normalized shear tensor. The parameter τ is a dimensionless time variable defined in terms of the Hubble scalar by

$$\frac{dt}{d\tau} = H^{-1}, \quad (9)$$

where t is the cosmological clock time. In an ever-expanding model, τ will assume all real values with

¹ An explicit introduction of a radiation fluid not interacting with the dust will not change the results.

$\tau \rightarrow +\infty$ at late times. The τ -dependence of Σ_{ab} and K^a is found by integrating the field equations and the null geodesic equations, which for spatially homogeneous models form a coupled system of ordinary differential equations (Nilsson *et al.* 1998). If the universe is *assumed* to be close to an FL model, then Σ_{ab} is small and Eq. (8) can be linearized in Σ_{ab} . Also, to lowest order, the directional cosines of the null geodesics in the corresponding background FL model can be used. For spatially homogeneous models, this approach has been used by various authors to obtain bounds on the Hubble normalized shear from CMB temperature observations. We refer, for example to Collins & Hawking (1973a), Barrow (1985), Bunn *et al.* (1996) and Kogut *et al.* (1997). Since we do *not* assume a priori that the models we are considering are close to an FL model, the full field equations and the exact null geodesic equations were integrated numerically to obtain the CMB temperature.

To quantify the anisotropy of the Bianchi type VII₀ dust models and hence their deviation from an FL model, we use the shear parameter Σ and the Weyl parameter $\mathcal{W}$ introduced previously. It has been shown by Collins & Hawking (1973b) that Bianchi VII₀ dust models initially close to the flat FL model isotropize at late times in the sense that $\Sigma \rightarrow 0$ as $\tau \rightarrow +\infty$. A more detailed analysis by Wainwright *et al.* (1998) showed that for any non-tilted Bianchi VII₀ dust model, the anisotropy parameters satisfy

$$\lim_{\tau \rightarrow +\infty} \Sigma = 0, \quad \lim_{\tau \rightarrow +\infty} \mathcal{W} = \mathcal{W}_\infty, \quad (10)$$

at late times. Here $\mathcal{W}_\infty$ is a constant whose value can be any positive number depending on the initial conditions. In other words, *although the models isotropize as regards the shear, the Weyl curvature remains dynamically significant*. This fact can be understood heuristically by considering the asymptotic form of the shear parameter Σ ,

$$\Sigma \approx C e^{-\frac{\tau}{2}} \left| \cos \left(\frac{4}{H_0 l_0} e^{\frac{\tau}{2}} + \psi_0 \right) \right|, \quad \tau \rightarrow +\infty, \quad (11)$$

where C, l_0, H_0 and ψ_0 are constants. This asymptotic form follows from Eq. (3.29), Eq. (3.33) and Eq. (C.10) in Wainwright *et al.* (1998). The decaying exponential ensures that Σ is small but the fact that Σ is highly oscillatory at late times generates a dynamically significant Weyl curvature.

We have performed a variety of numerical simulations to calculate anisotropy patterns of the CMB

temperature in Bianchi VII₀ models. Initial conditions are imposed at the present time τ_0 and then the field equations and null geodesics equations are integrated backwards to the time of last scattering τ_e . Then Eq. (8) is used to obtain the ratio T_o/T_e . The interval $\tau_0 - \tau_e$, corresponding to a redshift of $z \approx 1100$, is approximately 7. For a typical simulation, with present day values of the anisotropy parameters being $\Sigma_o = 3 \cdot 10^{-5}$ and $\mathcal{W}_o = 2$, we obtain a quadrupole of the order $a_2 \approx 7 \cdot 10^{-6}$ and an octupole of the order $a_3 \approx 7 \cdot 10^{-8}$. This model thus satisfies the observational constraint Eq. (6) but violates the theoretical upper bounds of Eq. (7) as regards the Weyl parameter. Hence, in the case of the present models, one or several of the hypotheses H_1, H_2, H_3 are not valid. Since the models we are considering are spatially homogeneous, hypothesis H_1 is valid and the calculations show that hypothesis H_2 is valid as well. It is thus, as expected, hypothesis H_3 that fails.

It is of interest to note that the octupole is considerably smaller than the quadrupole for these models. This is not a coincidence. Analytic considerations suggest that for Bianchi VII₀ models, a *pure* quadrupole pattern is obtained asymptotically at late times. Doroshkevich *et al.* (1975) have also given heuristic arguments that support this conclusion. It would be useful if this result could be proved and understood physically in a simple way since it is perhaps surprising. In order to investigate if it is possible to obtain detailed observationally compatible CMB temperature patterns using these models, one will have to consider density perturbations of these models. Will they give rise to similar CMB temperature maps as those obtained from density perturbations of FL models? Does a dynamically important Weyl curvature give rise to a different large scale structure and if so, does this in turn affect the CMB temperature?

In this letter we have shown that there exists cosmological models, specifically spatially homogeneous non-tilted Bianchi type VII₀ dust models, that are not close to any FL model even though the temperature of the CMB is almost isotropic in the sense that the observational bounds on the quadrupole and octupole are satisfied. It is also clear why this does not contradict the results of Maartens *et al.* (1995b) since one of the hypotheses on which their analysis is based is not valid for these models. Thus, the existence of these models suggests that hypothesis H_3 may not always be appropriate. It should be pointed out that the present class of models may very well be ruled out by

observations other than those of the CMB temperature. However, this is exactly the point. There is indeed a need for other cosmological observations.

We thank Roy Maartens for commenting on an earlier draft of the manuscript and Tim Gebbie for helpful discussions. This research was supported in part by a grant from the Natural Sciences & Engineering Research Council of Canada (JW), the Swedish Natural Research Council (CU), G  l  stiftelsen (USN), Svenska Institutet (USN), Stiftelsen Blanceflor (USN) and the University of Waterloo (USN, WCL).

REFERENCES

- Barrow, J. D., 1986, *Can. J. Phys.* 64, 152
- Bunn, E. F., Ferreira, P. G., Silk, J., 1996, *Phys. Rev. Lett.* 77, 2883
- Collins, C. B., Hawking, S. W., 1973a, *ApJ* 180, 317
- Collins, C. B., Hawking, S. W., 1973b, *MNRAS* 162, 307
- Doroshkevich, A.G., Lukash, V. N., Novikov, I. D., 1975, *Sov. Astron.* 18, 554
- Ehlers, J., Geren, P., Sachs, R. K., 1968, *J. Math. Phys.* 9, 1344
- Kogut, A., Hinshaw, G., Banday, A. J., 1997, *Phys. Rev. D* 55, 1901
- Lineweaver, C. H., Tenorjo, L., Smoot, G. F., Keegstra, P., Banday A. J., Lubin, P., 1996, *ApJ* 470, 38
- Kristian, J., Sachs, R. K., 1966, *ApJ* 143, 379
- Maartens, R., Ellis, G. F. R., Stoeger, W. R., 1995a, *Phys. Rev. D* 51, 1525
- Maartens, R., Ellis, G. F. R., Stoeger, W. R., 1995b, *Phys. Rev. D* 51, 5942
- Maartens, R., Ellis, G. F. R., Stoeger, W. R., 1996, *A&A* 309, L7
- Nilsson, U. S., Uggla, C., Wainwright, J., Lim, W. C., 1998, *in preparation*
- Stoeger, W. R., Maartens, R., Ellis, G. F. R., 1995, *ApJ* 443, 1
- Stoeger, W. R., Araujo, M. E., Gebbie, T., 1997, *ApJ* 476, 435
- Wainwright, J., Ellis, G. F. R., 1997, “Dynamical systems in cosmology”. Cambridge University Press, Cambridge
- Wainwright, J., Hancock, M., Uggla, C., 1998, “Asymptotic self-similarity breaking at late times in cosmology”, gr-qc/9812010, submitted to *Class. and Quantum Grav.*